Nudi

Large wound

SCIENCE FOR EVERY SPECIES

EQUINE CASE SERIES / 01

Charley McColough BVetMed, Whetstone Equine

PATIENT

16-year-old Quarter Horse gelding

OBSERVATION


40-day course

THE CASE

Following severe lymphangitis and cellulitis, the patient developed a large necrotic wound involving the right lateral hock and calcanean bursa. Initial care included systemic antibiotics and regional limb perfusion. Peptide therapy began after six days because of marked tissue loss and concern for communication with deep synovial structures.

REPORTED CARE

Daily subcutaneous BPC-157/TB-500 for 40 days, beginning February 23, 2026, alongside systemic antibiotics. Weekly bandage changes continued through March 9, 2026.



Right-hind wound: day 1 and day 40. Original photographs and date labels retained.

REPORTED OUTCOME

The clinician reported healthy, uniform granulation tissue, reduced discharge and progressive improvement in right-hind swelling. The patient remained comfortable. After 40 days, granulation tissue filled the cavity left by necrotic and infected tissue to the level of the surrounding skin.

Individual case report. Concurrent treatments were used; these observations do not establish efficacy or predict other patients' outcomes.

Nudi

EXPLORE THE CLINICAL LIBRARY

Scan for the research, case reports and resources behind the conversation.